

# Agilent Seahorse XF Media Selection Guide



Agilent Seahorse XF Media are specially formulated for use in XF Assays and are recommended for best results. XF Media are based on standard cell culture media composition (DMEM or RPMI), but varies from typical growth/culture medium formulations in a few key ways:

- **No bicarbonate and low buffering capacity:** to improve the detection of extracellular acidification.
- **No supplements:** allowing specific customization of the assay medium.
- **Low or no phenol red:** to allow for the most accurate and precise measurement of absolute pH values.

Agilent Seahorse now offers buffered XF media with pre-adjusted pH of 7.4 at 37 °C (part numbers 103575-100 and 103576-100). These media contain a low amount of HEPES buffer. When used with compatible XF supplements (e.g. XF glucose solution, XF pyruvate solution and XF glutamine solution) at recommended concentrations, there is no need to adjust pH of the media, thus, simplifying workflow and reducing time needed for assay preparation. These pH-ready media also have consistent buffering capacity, leading to more consistent assay data across experiments.

## Best practices for preparing Agilent Seahorse XF Assay Media using XF DMEM or RPMI media, pH 7.4

1. Transfer sufficient volume of XF DMEM or RPMI media, pH 7.4 to a new sterile bottle on the day of use.  
**Note:** It is recommended to not warm up the entire bottle of media if a smaller volume is needed, and to tighten the bottle cap after each use to maintain pH value.
2. Add desirable amounts of XF supplements/substrates.  
**Note:** XF supplements must be used at recommended concentration range to ensure a proper final pH in assay media. Proper media pH is not guaranteed if other brand supplements from other suppliers are used. Recommended supplement concentrations are 0 to 10 mM for glucose, 0 to 1 mM for pyruvate and 0 to 2 mM for glutamine. Specific supplement concentrations are assay dependent. See [Procedures for Preparing XF Assay Media](#) and [XF Assay Kit User Guides](#) for more information. Filter sterilization of the final assay media is not required if sterility of media and all supplements has not been compromised.
3. Warm medium to 37 °C. The assay medium is ready to use (no pH-adjustment is necessary).

## Best practices for preparing Agilent Seahorse XF Assay Media using media that require pH adjustment

1. Transfer a sufficient volume of XF medium to a new bottle on the day of use.
2. Add desirable amounts of supplements/substrates.  
**Note:** For preparing assay media to use in Glycolytic Rate Assay, HEPES buffer must be added at 5 mM for the DMEM medium composition and 1 mM for RPMI medium composition. For media used in other XF assays, inclusion of HEPES is optional.
3. Warm medium to 37 °C.
4. Adjust pH to 7.4 ±0.1.
5. Filter-sterilize the medium after adjusting pH value. The assay medium is ready to use.

## Agilent Seahorse XF Media, Buffer, and supplement products

Part No.	Product Name	Size	Core Formula	Phenol Red	Note
103575-100	Seahorse XF DMEM Medium, pH 7.4	500 mL	DMEM	No	Suitable for all XF assays. Contains HEPES.
103576-100	Seahorse XF RPMI Medium, pH 7.4	500 mL	RPMI	No	Suitable for all XF assays. Contains HEPES.
103335-100	Seahorse XF Base Medium (without Phenol Red)	500 mL	DMEM	No	Suitable for all XF assays. 5 mM HEPES is required for Glycolytic Rate Assay.
102353-100	Seahorse XF Base Medium	2 × 1 L	DMEM	Yes	Suitable for all XF assays except for Glycolytic Rate Assay.
103334-100	Seahorse XF Base Medium, 500 mL	500 mL	DMEM	Yes	Suitable for all XF assays except for Glycolytic Rate Assay.
103193-100	Seahorse XF Base Medium, 100 mL	100 mL	DMEM	Yes	Suitable for all XF assays except for Glycolytic Rate Assay.
103577-100	Seahorse XF 1.0 M Glucose Solution, 50 mL	50 mL	n/a	No	Compatible with all XF media.
103578-100	Seahorse XF 100 mM Pyruvate Solution, 50 mL	50 mL	n/a	No	Compatible with all XF media.
103579-100	Seahorse XF 200 mM Glutamine Solution, 50 mL	50 mL	n/a	No	Compatible with all XF media.

**Note:** All Seahorse XF Media/Buffer/Supplements are endotoxin tested and should be stored at 4 °C, except for glutamine solution which should be stored at -20 °C

## Agilent Seahorse XF Media, Buffer, and supplement products

Part No.	Product Name	Real-Time ATP Rate Assay	Cell Mito Stress Test	Glycolytic Rate Assay	Glycolysis Stress Test	Cell Energy Phenotype Test	Mito Fuel Flex Test	Substrate Oxidation Stress Test	Palmitate Oxidation Stress Test
103575-100	Seahorse XF DMEM Medium, pH 7.4	●	●	●	●	●	●	●	●
103576-100	Seahorse XF RPMI Medium, pH 7.4	✓	✓	✓	✓	✓	✓	✓	✓
103335-100	Seahorse XF Base Medium (without Phenol Red)	✓●	✓	✓●	✓	✓	✓	✓	✓
102353-100	Seahorse XF Base Medium, 2 × 1 L	●	✓	●	✓	✓	✓	✓	✓
103334-100	Seahorse XF Base Medium, 500 mL	●	✓	●	✓	✓	✓	✓	✓
103193-100	Seahorse XF Base Medium, 100 mL	●	✓	●	✓	✓	✓	✓	✓

- Recommended
- ✓ Compatible
- Not Compatible
- Requires addition of HEPES

### Your Contact in Switzerland:



**Bucher Biotec AG**  
 Viaduktstrasse 42  
 CH-4051 Basel  
 Tel. 061 269 1111  
[info@bucher.ch](mailto:info@bucher.ch)  
[www.bucher.ch](http://www.bucher.ch)



[www.agilent.com/chem/discoverxf](http://www.agilent.com/chem/discoverxf)

**For Research Use Only. Not for use in diagnostic procedures.**

This information is subject to change without notice.

© Agilent Technologies, Inc. 2020  
 Printed in the USA, February 3, 2020  
 5994-7878EN  
 DE.3245601852

Bucher Biotec AG is the official channel partner of Agilent for all Seahorse products in Switzerland.

- Reagents, Consumables and Instruments -

For your own convenience and fastest delivery times, please place your orders directly with Bucher Biotec AG

[orders@bucher.ch](mailto:orders@bucher.ch)